

# Reciprocal and Variational Principles in Micropolar Thermoelasticity \*

*Francesca Passarella*

Dipartimento di Ingegneria dell'Informazione e Matematica Applicata  
Università di Salerno - via Ponte don Melillo - I-84084 Fisciano (SA)

---

\*Work performed in the context of a fellowship of Italian Research Council (C.N.R.).

## References

- [1] A. C. Eringen: *Linear Theory of Micropolar Elasticity*. Int. Math. Mech. **15**, 909 (1966).
- [2] A. C. Eringen: *Fondamentions of Micropolar Thermoelasticity*. Springer–Verlag, Vienna (1970).
- [3] A. C. Eringen: *Theory of Micropolar elasticity*. In Fracture (Edited by H. Leibowitz), Vol II, Academic Press, New York, 622 (1968).
- [4] A. C. Eringen and C. B. Kafadar: *Polar Field Theories*. In Continuum Phisics (Edited by A. C. Eringen), Vol IV, Part I, Academic Press, New York, 1 (1976).
- [5] M. Ciarletta and D. Ieşan: *Non–classical elastic solids*. In Pitman Research Notes in Mathematics Series (Edited by Longman Scientific & Technical), 293 (1993).
- [6] D. S. Chandrasekharaiah: *Heat–flux dependent micropolar thermoelasticity*, Int. J. Engng. Sci. **24**, 1389 (1986).
- [7] M. Ciarletta: *Sui processi termoelastici per continui micropolari*, Atti Sem. Mat. Fis. Univ. Modena, **39**, 103 (1991).
- [8] J. W. Nunziato and S. C. Cowin: *A nonlinear theory of elastic materials with voids*, Arch. Rat. Mech. Anal. **72**, 175 (1979).
- [9] S. C. Cowin and J. W. Nunziato: *Linear elastic materials with voids*, J. Elasticity **13**, 125 (1983).
- [10] M. A. Goodman and S. C. Cowin: *A continuum theory for granular materials*, Arch. Rat. Mech. Anal. **44**, 249 (1972).
- [11] F. Passarella: *Some results in micropolar thermoelasticity*, pre-print n.10/95 D.I.I.M.A., University of Salerno.
- [12] M.A. Biot: *Thermoelasticity and irreversible thermodynamics*, J. Appl. Phys. **7**, 240 (1956)
- [13] D. S. Chandrasekharaiah: *Variational and reciprocal principles in micropolar thermoelasticity*, Int. J. Engng. Sci. **25**, 55 (1987).

# Some theorems in thermoelasticity for micropolar porous media

*Gerardo Iovane*

*Francesca Passarella*

Dipartimento di Ingegneria dell'Informazione e Matematica Applicata  
Università di Salerno - via Ponte don Melillo - I-84084 Fisciano (SA)

## Abstract

Within the context of a linear theory of heat-flux dependent thermoelasticity for micropolar porous media some variational principles and a reciprocal relation are derived.

## 1 Introduction

This paper is concerned with the linear theory of the generalized thermodynamics for the micropolar porous media.

The *micropolar elasticity* theory has been established by Eringen [1-4] and is of fundamental interest in a lot of cases for which the classical theory of elasticity is inadequate. Other related results have been obtained in [5-7].

A general theory of materials with voids has been developed by Cowin and Nunziato in [8-9], by using the concept of *distributed body* introduced in [10]. In this theory, the matrix material is elastic and the interstices are void of material. In this context, the *bulk density* is written as the product of two fields: the material density field and the volume fraction field. The intended applications of this theory are to geological materials (like rock and soils) and to manufactured porous materials (like ceramics and pressed powders).

In a previous paper [11], we studied the thermodynamic theory of micropolar porous materials by using the theories developed in [1-4], [6] and in [8-9]. This study was based on a generalized theory of thermoelasticity including the *heat flux vector* in the set of the constitutive variables. In [11], we also derived the corresponding linear theory.

In the present work, starting from the basic equations established in [11], we state two variational characterization (of Biot and Hamilton types, see [12], [13]) of the initial-boundary value problem. In final part, we also derive a reciprocal relation for the problem in concern by using convolution in time (fundamental results, in this context, have been obtained by D. Graffi [14]) and Laplace transform.

## 2 Basic Equations

Throughout this paper, by  $\Omega$  and  $\partial\Omega$  we shall denote the region and its boundary of the physical 3-dimensional space ( $\equiv \mathbb{R}^3$ ) occupied by an elastic solid in a given reference (unstressed) state and  $I = [0, +\infty)$ . We refer the motion of the body to a fixed orthonormal frame in  $\mathbb{R}^3$ . We shall denote the tensor components of order  $p \geq 1$  by Latin subscripts, ranging over  $\{1, 2, 3\}$ . Summation over repeated subscripts is implied. Superposed dots or subscripts preceded by a comma will mean partial derivative with respect to the time or the corresponding coordinates.

On the basis of the linear theory established in [11], the behaviour of a micropolar porous thermoelastic body is governed by the following local balances of momentum, equilibrated force and energy

$$T_{ji,j} + \rho f_i = \rho \ddot{u}_i, \tag{1}$$

$$M_{ji,j} + \epsilon_{irs} T_{rs} + \rho \ell_i = \rho J_{ij} \ddot{\phi}_j$$

$$h_{i,i} + g + \rho \ell = \rho \chi \ddot{\varphi},$$

$$\rho(\theta_0 \dot{\eta} - r) = q_{i,i},$$

where

$T_{ij}, M_{ij}$  stress and couple stress tensors;  
 $h_i, g$  equilibrated stress vector, intrinsic equilibrated body force;  
 $\eta, q_i$  entropy and heat flux vector.

These are the dependent variables of the theory, needing a constitutive equation.

Further,  $\rho$  is the bulk mass density,  $f_i$  the body force,  $\ell_i$  the body couple,  $J_{ij}$  the micro inertia tensor,  $\chi$  the equilibrated inertia,  $\ell$  the extrinsic equilibrated body force,  $r$  the strength of the internal heat source and  $\theta_0$  the (constant  $> 0$ ) temperature in the reference state;  $\epsilon_{irs}$  is the alternating symbol.

Finally,  $u_i, \phi_i$  and  $\varphi$ , along with the temperature  $\theta$  (measured from  $\theta_0$ ), are the thermokinetic variables of the theory:  $u_i$  represents the displacement,  $\phi_i$  the micro-rotation,  $\varphi$  the change in volume fraction from the reference configuration.

The constitutive equations for (1) are [11]

$$\begin{aligned}
T_{ij} &= C_{ijkl}E_{kl} + G_{ijkl}\Psi_{lk} + H_{ij}\varphi + H_{ijk}\varphi_{,k} + A_{ij}\theta, \\
M_{ij} &= G_{klij}E_{kl} + \Gamma_{ijkl}\Psi_{lk} + P_{ij}\varphi + P_{ijk}\varphi_{,k} + G_{ij}\theta, \\
g &= -H_{ij}E_{ij} - P_{ij}\Psi_{ji} - a\varphi - a_i\varphi_{,i} - b\theta, \\
h_i &= H_{jki}E_{jk} + P_{jki}\Psi_{kj} + a_i\varphi + D_{ij}\varphi_{,j} + \gamma_i\theta, \\
\rho\eta &= -A_{ij}E_{ij} - G_{ij}\Psi_{ji} - b\varphi - \gamma_i\varphi_{,i} + c\theta, \\
(1 + \tau\frac{\partial}{\partial t})q_i &= K_{ij}\theta_{,j}.
\end{aligned} \tag{2}$$

Here,  $E_{ij}$  and  $\Psi_{ij}$  are the *deformation* and *wryness tensors* defined by

$$E_{ij} = u_{j,i} - \epsilon_{ijh}\phi_h, \quad \Psi_{ij} = \phi_{i,j}. \tag{3}$$

We shall consider a homogeneous body; in this case, the coefficients in (2) are all constants; it can also be proved [11] that the following symmetry relations follow

$$C_{ijkl} = C_{klij}, \quad \Gamma_{ijkl} = \Gamma_{klij} \quad D_{ij} = D_{ji} \quad (4)$$

and that the conductivity tensor  $K_{ij}$  is inversable.

Note that applying the operator  $\left(1 + \tau \frac{\partial}{\partial t}\right)$  to the energy equation, by (2<sub>5,6</sub>) we get an evolution equation for  $\theta$  of hyperbolic type

$$K_{ij}\theta_{,ji} + \left(1 + \tau \frac{\partial}{\partial t}\right) [\rho r - \rho \theta_0 \dot{\eta}] = 0. \quad (5)$$

By insertion of (2, 3) into (1<sub>1,2,3</sub>, 5), we obtain the complete system of (coupled) field equations for  $u_i$ ,  $\phi_i$ ,  $\varphi$  and  $\theta$ .

Let us denote four pairs of disjoint and complementary subsets of the (smooth) boundary  $\partial\Omega$  by  $\{\Sigma_i, \Sigma_{i+1}\}$  with  $i = 1, 3, 5, 7$  and denote the outward unit normal to  $\partial\Omega$  by  $n_j$ .

To the system of field equations (1<sub>1,2,3</sub>, 2, 5) we add the following initial–boundary conditions

$$u_i = \dot{u}_i = \phi_i = \dot{\phi}_i = \varphi = \dot{\varphi} = \theta = q_i = 0 \quad \text{in } \Omega \times \{0\}, \quad (6)$$

$$\begin{aligned} u_i &= u_i^* & \text{on } \Sigma_1 \times I, & & T_{ji}n_j &= t_i^* & \text{on } \Sigma_2 \times I, \\ \phi_i &= \phi_i^* & \text{on } \Sigma_3 \times I, & & M_{ji}n_j &= m_i^* & \text{on } \Sigma_4 \times I, \\ \varphi &= \varphi^* & \text{on } \Sigma_5 \times I, & & h_j n_j &= h^* & \text{on } \Sigma_6 \times I, \\ \theta &= \theta^* & \text{on } \Sigma_7 \times I, & & q_j n_j &= q^* & \text{on } \Sigma_8 \times I, \end{aligned} \quad (7)$$

where right-hand terms stand for (sufficiently smooth) assigned fields; along with  $f_i$ ,  $\ell_i$ ,  $\ell$  and  $r$ , these are the *data* of the mixed problem considered. An array field  $(u_i, \phi_i, \varphi, \theta)$  meeting all equations (1, 2, 3, 5, 6, 7), for some assignment of the data, will be referred to as a (regular) *solution* of this.

The uniqueness of solution to the above problem has been established in [11].

### 3 Variational principle I

In this section, following [15], we establish a variational theorem of Hamilton type. We introduce (see also [11]) the *kinetic energy*  $K$  and the *free energy*  $\psi$

$$\begin{aligned} \rho K = & \frac{1}{2}\rho(\dot{u}_i\dot{u}_i + J_{ij}\dot{\phi}_i\dot{\phi}_j + \chi\dot{\varphi}^2) \\ \rho\psi = & \frac{1}{2}C_{ijkl}E_{ij}E_{kl} + \frac{1}{2}\Gamma_{ijkl}\Psi_{ji}\Psi_{lk} + \frac{1}{2}a\varphi^2 + \frac{1}{2}D_{ij}\varphi_{,i}\varphi_{,j} - \frac{1}{2}c\theta^2 + \\ & + \frac{1}{2}B_{ij}q_iq_j + G_{ijkl}E_{ij}\Psi_{lk} + H_{ij}E_{ij}\varphi + H_{ijk}E_{ij}\varphi_{,k} + A_{ij}E_{ij}\theta + \\ & + P_{ij}\Psi_{ji}\varphi + P_{ijk}\Psi_{ji}\varphi_{,k} + G_{ij}\Psi_{ji}\theta + a_i\varphi\varphi_{,i} + b\varphi\theta + \gamma_i\varphi_{,i}\theta. \end{aligned} \tag{8}$$

We can prove the following

**Variational principle I** *Let  $t_1$  and  $t_2$  be arbitrary instants of time in  $I$  and assume that*

- i. the symmetry relations (4) hold;*
- ii.  $K_{ij}$  and  $J_{ij}$  are symmetric tensors.*

*Then, for arbitrary variations in  $u_i$ ,  $\phi_i$ ,  $\varphi$ ,  $\theta$  such that*

$$\delta u_i = 0, \quad \delta \phi_i = 0, \quad \delta \varphi = 0, \quad \delta \theta = 0, \quad \text{in } \Omega \times \{t_1, t_2\}, \tag{9}$$

$$\begin{aligned}
\delta u_i &= 0 & \text{on } \Sigma_1 \times I, & \delta \phi_i &= 0 & \text{on } \Sigma_3 \times I, \\
\delta \varphi &= 0 & \text{on } \Sigma_5 \times I, & \delta \theta &= 0 & \text{on } \Sigma_7 \times I,
\end{aligned} \tag{10}$$

and the functions  $f_i, \ell_i, \ell, \eta, r, t_i^*, m_i^*, h^*, q^*, t_1$  and  $t_2$  being kept unchanged, the following variational equations

$$\begin{aligned}
\delta \int_{t_1}^{t_2} dt \left[ \int_{\Omega} \rho \left( K - \psi - \eta T + \frac{1}{2\rho} B_{ij} q_i q_j + f_i u_i + \ell_i \phi_i + \ell \varphi \right) dV + \right. \\
\left. + \int_{\Sigma_2} t_i^* u_i d\sigma + \int_{\Sigma_4} m_i^* \phi_i d\sigma + \int_{\Sigma_6} h^* \varphi d\sigma \right] = 0
\end{aligned} \tag{11}$$

$$\delta \int_{t_1}^{t_2} dt \left[ \int_{\Omega} \frac{1}{2} K_{ij} T_{,i} T_{,j} dV - (1 + \tau p) \left\{ \int_{\Omega} \rho (\eta \dot{T} - r) T dV + \int_{\Sigma_8} q^* T d\sigma \right\} \right] = 0 \tag{12}$$

are satisfied if and only if are verified the field equations  $(1_{1,2,3})$  and  $(5)$ . In the above equations we put  $T = \theta + \theta_0$  (the absolute temperature) and  $p \equiv \frac{\partial}{\partial t}$ .

**Proof.** We can remark that, under the conditions (9), the variation of the kinetic energy is

$$\delta \int_{t_1}^{t_2} dt \int_{\Omega} \rho K dV = - \int_{t_1}^{t_2} dt \int_{\Omega} \rho \left( \ddot{u}_i \delta u_i + J_{ij} \ddot{\phi}_j \delta \phi_i + \chi \ddot{\varphi} \delta \varphi \right) dV. \tag{13}$$

Moreover, using  $(8_2, 2, 3, 4, 7, 9, 10)$  and the divergence theorem, we can obtain



$$\begin{aligned}
& \delta \int_{t_1}^{t_2} dt \int_{\Omega} \left( \rho \psi + \rho \eta T - \frac{1}{2} B_{ij} q_i q_j \right) dV = \\
& = \int_{t_1}^{t_2} dt \int_{\Omega} (T_{ji} \delta E_{ji} + M_{ji} \delta \Psi_{ij} - g \delta \varphi + h_i \delta \varphi_{,i}) dV = \\
& = \int_{t_1}^{t_2} dt \left[ \int_{\Omega} \left\{ -T_{ji,j} \delta u_i - (M_{ji,j} + \epsilon_{irs} T_{rs}) \delta \phi_i - (h_{i,i} + g) \delta \varphi \right\} dV + \right. \\
& \quad \left. + \int_{\Sigma_2} t_i^* \delta u_i d\sigma + \int_{\Sigma_4} m_i^* \delta \phi_i d\sigma + \int_{\Sigma_6} h^* \delta \varphi d\sigma \right].
\end{aligned} \tag{14}$$

Taking into account (13) and (14), the variational equation (11) can be rewritten as follows

$$\begin{aligned}
& \int_{t_1}^{t_2} dt \int_{\Omega} \left[ (T_{ji,j} + \rho f_i - \rho \ddot{u}_i) \delta u_i + (M_{ji,j} + \epsilon_{irs} T_{rs} + \rho \ell_i - \rho J_{ij} \ddot{\phi}_j) \delta \phi_i \right. \\
& \quad \left. + (h_{i,i} + g + \rho \ell - \rho \chi \ddot{\varphi}) \delta \varphi \right] dV = 0
\end{aligned}$$

This equation holds if and only if (1<sub>1,2,3</sub>) are satisfied.

From (2<sub>6</sub>) and (10) it follows that

$$\delta \int_{t_1}^{t_2} dt \int_{\Omega} \frac{1}{2} K_{ij} T_{,i} T_{,j} dV = \int_{t_1}^{t_2} dt \left[ (1 + \tau p) \int_{\Sigma_8} q^* \delta \theta d\sigma - \int_{\Omega} K_{ij} \theta_{,ij} \delta \theta dV \right]$$

where  $p$  has been treated as a constant [12].

Thus, (12) can be written as

$$\int_{t_1}^{t_2} dt \int_{\Omega} \{ K_{ij} \theta_{,ij} + \rho(1 + \tau p)(r - T \dot{\eta}) \} \delta \theta dV = - \int_{\Omega} (\eta T \delta T)|_{t_1}^{t_2} dV$$

After the linearization, from (9) we conclude that

$$\int_{t_1}^{t_2} dt \int_{\Omega} \{ K_{ij} \theta_{,ij} + \rho(1 + \tau p)(r - \theta_0 \dot{\eta}) \} \delta \theta dV = 0$$

This equation holds if and only if (5) is satisfied.

## 4 Variational pinciple II

Following [12], we introduce the *entropy flow* vector  $s_i$  by

$$\theta_0 \dot{s}_i = q_i$$

and by (1<sub>4</sub>, 2<sub>5</sub>) we get

$$\dot{s}_{i,i} = -A_{ij}\dot{E}_{ij} - G_{ij}\dot{\Psi}_{ji} - b\dot{\varphi} - \gamma_i\dot{\varphi}_{,i} + c\dot{\theta} - \frac{\rho}{\theta_0}r. \quad (15)$$

In the following, we shall consider prescribed  $f_i$ ,  $\ell_i$ ,  $\ell$  and  $r$  in  $\Omega$ , and  $u_i$  on  $\Sigma_1$ ,  $T_{ij}$  on  $\Sigma_2$ ,  $\phi_i$  on  $\Sigma_3$ ,  $M_{ij}$  on  $\Sigma_4$ ,  $\varphi$  on  $\Sigma_5$ ,  $h_i$  on  $\Sigma_6$ ,  $\theta$  on  $\Sigma_7$  and  $q_i$  on  $\Sigma_8$ . Moreover, we also consider  $s_i$  assigned on  $\Sigma_8$ .

If we assume that  $K_{ij}$  is symmetric and denote its inverse by  $\widetilde{K}_{ij}(= \widetilde{K}_{ji})$ , we set

$$D = \frac{1}{2} \int_{\Omega} \theta_0(p + \tau p^2) \widetilde{K}_{ij} s_i s_j dV$$

so the variation of this functional due to variation of  $s_i$  is

$$\delta D = \int_{\Omega} \theta_0(p + \tau p^2) \widetilde{K}_{ij} s_i \delta s_j dV. \quad (16)$$

Here,  $p$  is used as in section 3.

We now introduce the following functional

$$V = \int_{\Omega} (\rho\psi + \rho\eta\theta - \frac{1}{2}B_{ij}q_iq_j) dV.$$

If we consider variations of  $V$  due to variations of  $u_i$ ,  $\phi_i$ ,  $\varphi$ ,  $\theta$  and use (8<sub>2</sub>), the

constitutive equations and the symmetry relations (4), we obtain

$$\begin{aligned} \delta V = \int_{\Omega} \{ & c\theta\delta\theta + (T_{ij} - A_{ij}\theta)\delta E_{ij} + (M_{ij} - G_{ij}\theta)\delta\Psi_{ji} + \\ & -(g + b\theta)\delta\varphi + (h_i - \gamma_i\theta)\delta\varphi_{,i} \} dV. \end{aligned} \quad (17)$$

Moreover, we define the following functional

$$G = - \int_{\Sigma_2} t_i^* u_i d\sigma - \int_{\Sigma_4} m_i^* \phi_i d\sigma - \int_{\Sigma_6} h^* \varphi d\sigma - \int_{\Sigma_7} \theta^* s_i n_i d\sigma$$

If  $\delta u_i$ ,  $\delta\phi_i$ ,  $\delta\varphi$ ,  $\delta s_i$  vanish on  $\Sigma_1$ ,  $\Sigma_3$ ,  $\Sigma_5$ ,  $\Sigma_8$ , respectively, by (7) we can write

$$\delta G = - \int_{\partial\Omega} (T_{ji}\delta u_i + M_{ji}\delta\phi_i + h_j\delta\varphi + \theta\delta s_j) n_j d\sigma;$$

and, from the divergence theorem and (3, 4, 15), we get

$$\begin{aligned} \delta G = - \int_{\Omega} \{ & (T_{ij} - A_{ij}\theta)\delta E_{ij} + (M_{ij} - G_{ij}\theta)\delta\Psi_{ji} + (h_i - \gamma_i\theta)\delta\varphi_{,i} + c\theta\delta\theta + \\ & -b\theta\delta\varphi + T_{ji,j}\delta u_i + (M_{ji,j} + \epsilon_{irs}T_{rs})\delta\phi_i + h_{j,j}\delta\varphi + \theta_{,j}\delta s_j \} dV. \end{aligned} \quad (18)$$

Finally, we put

$$F = \int_{\Omega} \rho \left\{ \left( \frac{1}{2} p^2 u_i - f_i \right) u_i + \left( \frac{1}{2} p^2 J_{ij} \phi_j - \ell_i \right) \phi_i + \left( \frac{1}{2} \chi p^2 \varphi - \ell \right) \varphi \right\} dV,$$

so, assuming the tensor  $J_{ij}$  symmetric, the variation of this functional is

$$\delta F = \int_{\Omega} \rho \left\{ (p^2 u_i - f_i) \delta u_i + (p^2 J_{ij} \phi_j - \ell_i) \delta \phi_i + (\chi p^2 \varphi - \ell) \delta \varphi \right\} dV. \quad (19)$$

Thus, we can obtain the following variational theorem (of the Biot type)

**Variational pinciple II** *Let us prescribe  $f_i, \ell_i, \ell$  and  $r$  in  $\Omega$ , and  $T_{ij}, M_{ij}, h_i$ , and  $\theta$  on  $\Sigma_2, \Sigma_4, \Sigma_6, \Sigma_7$ , respectively, and assume that*

- i. the symmetry relations (4) hold;*
- ii.  $K_{ij}$  and  $J_{ij}$  are symmetric tensors.*

*Then, for every  $\delta u_i$  ( $= 0$  on  $\Sigma_1$ ),  $\delta \phi_i$  ( $= 0$  on  $\Sigma_3$ ),  $\delta \varphi$  ( $= 0$  on  $\Sigma_5$ ),  $\delta s_i$  ( $= 0$  on  $\Sigma_8$ ), the following equation*

$$\delta H(u_i, \phi_i, \varphi, \theta) = \delta V + \delta G + \delta F + \delta D = 0$$

*yields the field equations (1<sub>1,2,3</sub>, 5).*

**Proof.** By (16, 17, 18, 19) we can write

$$\begin{aligned} \delta H(u_i, \phi_i, \varphi, \theta) = & \int_{\Omega} \left\{ (\rho \ddot{u}_i - \rho f_i - T_{ji,j}) \delta u_i + \right. \\ & + (\rho J_{ij} \ddot{\phi}_i - \rho \ell_i - M_{ji,j} - \epsilon_{irs} T_{rs}) \delta \phi_i + \\ & + (\rho \chi \ddot{\varphi} - \rho \ell - g - h_{i,i}) \delta \varphi + \\ & \left. + [\theta_0(1 + \tau p) \widetilde{K}_{ij} \dot{s}_i - \theta_{,j}] \delta s_i \right\} dV. \end{aligned}$$

Vanishing of this expression for arbitrary  $\delta u_i, \delta \phi_i, \delta \varphi, \delta s_i$  of course implies the field equations 1<sub>1,2,3</sub>; further from the last term in the integral above, we get

$$\theta_0(1 + \tau p) \widetilde{K}_{ij} \dot{s}_i = \theta_{,j}.$$

If we multiply the above equation by  $K_{hi}$  and derive with respect to the  $i$ -th coordinate, we can easily deduce the field equation (5) from (15). This completes the proof.

## 5 Reciprocity

Let us denote by  $a * b$  the time convolution of the scalar fields on  $\Omega \times I$

$$(a_1 * a_2)(\mathbf{x}, t) = \int_0^t a_1(\mathbf{x}, t - \tau) a_2(\mathbf{x}, \tau) d\tau.$$

Let us consider the body subjected to two different sets of data

$$\mathcal{T}^{(\alpha)} = \left\{ f_i^{(\alpha)}, \ell_i^{(\alpha)}, \ell^{(\alpha)}, r^{(\alpha)}, u_i^{*(\alpha)}, t_i^{*(\alpha)}, \phi_i^{*(\alpha)}, m_i^{*(\alpha)}, \varphi^{*(\alpha)}, h^{*(\alpha)}, \theta^{*(\alpha)}, q^{*(\alpha)} \right\}$$

and null initial conditions and let be

$$\mathcal{U}^{(\alpha)} = \left( u_i^{(\alpha)}, \phi_i^{(\alpha)}, \varphi^{(\alpha)}, \theta^{(\alpha)} \right) \quad \alpha = 1, 2,$$

the corresponding solutions. Moreover, we define  $T_{ij}^{(\alpha)}, M_{ij}^{(\alpha)}, h_i^{(\alpha)}, g^{(\alpha)}, \eta^{(\alpha)}, q_i^{(\alpha)}$  by means of (2) for each  $\alpha = 1, 2$ .

We prove the following theorem

**Reciprocal Theorem** *Let  $\mathcal{U}^{(\alpha)}$  be solutions corresponding to different sets of data  $\mathcal{T}^{(\alpha)}$  ( $\alpha = 1, 2$ ), and assume that*

- i. the symmetry relations (4) hold;*
- ii.  $K_{ij}$  and  $J_{ij}$  are symmetric tensors.*

*Then, the following relation holds*

$$\mathcal{I}_{\alpha\beta} = \mathcal{I}_{\beta\alpha} \tag{20}$$

where

$$\begin{aligned}
\mathcal{I}_{\alpha\beta} = & \int_{\Omega} \rho \left[ f_i^{(\alpha)} * \dot{u}_i^{(\beta)} + \ell_i^{(\alpha)} * \dot{\phi}_i^{(\beta)} + \right. \\
& \left. + \ell^{(\alpha)} * \dot{\varphi}^{(\beta)} - \frac{1}{\theta_0} r^{(\alpha)} * \theta^{(\beta)} \right] dV + \\
& - \int_{\Sigma_1} \dot{u}_i^{*(\alpha)} * T_{ji}^{(\beta)} n_j d\sigma + \int_{\Sigma_2} t_i^{*(\alpha)} * \dot{u}_i^{(\beta)} d\sigma + \\
& - \int_{\Sigma_3} \dot{\phi}_i^{*(\alpha)} * M_{ji}^{(\beta)} n_j d\sigma + \int_{\Sigma_4} m_i^{*(\alpha)} * \dot{\phi}_i^{(\beta)} d\sigma + \\
& - \int_{\Sigma_5} \dot{\varphi}^{*(\alpha)} * h_j^{(\beta)} n_j d\sigma + \int_{\Sigma_6} h^{*(\alpha)} * \dot{\varphi}^{(\beta)} d\sigma + \\
& + \frac{1}{\theta_0} \int_{\Sigma_7} \theta^{*(\alpha)} * q_j^{(\beta)} n_j d\sigma - \frac{1}{\theta_0} \int_{\Sigma_8} q^{*(\alpha)} * \theta^{(\beta)} d\sigma
\end{aligned}$$

for  $\alpha, \beta = 1, 2$ .

**Proof.** We put

$$\begin{aligned}
\mathcal{F}_{\alpha\beta} = & \left( \tilde{T}_{ji}^{(\alpha)} \tilde{u}_i^{(\beta)} + \tilde{M}_{ji}^{(\alpha)} \tilde{\phi}_i^{(\beta)} + \tilde{h}_j^{(\alpha)} \tilde{\varphi}^{(\beta)} \right)_{,j} + \\
& + \rho \left( \tilde{f}_i^{(\alpha)} \tilde{u}_i^{(\beta)} + \tilde{l}_i^{(\alpha)} \tilde{\phi}_i^{(\beta)} + \tilde{\ell}^{(\alpha)} \tilde{\varphi}^{(\beta)} \right), \\
\mathcal{H}_{\alpha\beta} = & \left( \tilde{\theta}^{(\alpha)} \tilde{q}_i^{(\beta)} \right)_{,i} - \rho \tilde{r}^{(\alpha)} \tilde{\theta}^{(\beta)}
\end{aligned}$$

where  $\tilde{a}$  is Laplace transform with respect to  $t$  of the function  $a$ :  $\tilde{a}(s) = \int_0^\infty e^{-st} a(t) dt$ . If we take Laplace transform of (1, 2) under the hypotheses  $i$ ,  $ii$  and the initial conditions (6), we deduce that

$$\begin{aligned}
\mathcal{F}_{\alpha\beta} - \mathcal{F}_{\beta\alpha} = & A_{ij}(\tilde{\theta}^{(\alpha)}\tilde{E}_{ij}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{E}_{ij}^{(\alpha)}) + G_{ij}(\tilde{\theta}^{(\alpha)}\tilde{\Psi}_{ji}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{\Psi}_{ji}^{(\alpha)}) + \\
& + b(\tilde{\theta}^{(\alpha)}\tilde{\varphi}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{\varphi}^{(\alpha)}) + \gamma_i(\tilde{\theta}^{(\alpha)}\tilde{\varphi}_{,i}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{\varphi}_{,i}^{(\alpha)}) \\
\mathcal{H}_{\alpha\beta} - \mathcal{H}_{\beta\alpha} = & -\theta_0 s \left[ A_{ij}(\tilde{\theta}^{(\alpha)}\tilde{E}_{ij}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{E}_{ij}^{(\alpha)}) + G_{ij}(\tilde{\theta}^{(\alpha)}\tilde{\Psi}_{ji}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{\Psi}_{ji}^{(\alpha)}) + \right. \\
& \left. + b(\tilde{\theta}^{(\alpha)}\tilde{\varphi}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{\varphi}^{(\alpha)}) + \gamma_i(\tilde{\theta}^{(\alpha)}\tilde{\varphi}_{,i}^{(\beta)} - \tilde{\theta}^{(\beta)}\tilde{\varphi}_{,i}^{(\alpha)}) \right],
\end{aligned}$$

so

$$s(\mathcal{F}_{\alpha\beta} - \mathcal{F}_{\beta\alpha}) + \frac{1}{\theta_0}(\mathcal{H}_{\alpha\beta} - \mathcal{H}_{\beta\alpha}) = 0 \quad (21)$$

If we integrate this equation over  $\Omega$  and use the divergence theorem and boundary conditions, we get

$$\begin{aligned}
& \int_{\Omega} \rho s \left[ \tilde{f}_i^{(\alpha)}\tilde{u}_i^{(\beta)} + \tilde{l}_i^{(\alpha)}\tilde{\phi}_i^{(\beta)} + \tilde{\ell}^{(\alpha)}\tilde{\varphi}^{(\beta)} - \tilde{f}_i^{(\beta)}\tilde{u}_i^{(\alpha)} + \right. \\
& \quad \left. - \tilde{l}_i^{(\beta)}\tilde{\phi}_i^{(\alpha)} - \tilde{\ell}^{(\beta)}\tilde{\varphi}^{(\alpha)} \right] - \frac{\rho}{\theta_0} \left[ \tilde{r}^{(\alpha)}\tilde{\theta}^{(\beta)} - \tilde{r}^{(\beta)}\tilde{\theta}^{(\alpha)} \right] dV + \\
& + \int_{\Sigma_1} s(\tilde{T}_{ji}^{(\alpha)}\tilde{u}_i^{*(\beta)} - \tilde{T}_{ji}^{(\beta)}\tilde{u}_i^{*(\alpha)})n_j d\sigma + \int_{\Sigma_2} s(\tilde{t}_i^{*(\alpha)}\tilde{u}_i^{(\beta)} - \tilde{t}_i^{*(\beta)}\tilde{u}_i^{(\alpha)}) d\sigma + \\
& + \int_{\Sigma_3} s(\tilde{M}_{ji}^{(\alpha)}\tilde{\phi}_i^{*(\beta)} - \tilde{M}_{ji}^{(\beta)}\tilde{\phi}_i^{*(\alpha)})n_j d\sigma + \int_{\Sigma_4} s(\tilde{m}_i^{*(\alpha)}\tilde{\phi}_i^{(\beta)} - \tilde{m}_i^{*(\beta)}\tilde{\phi}_i^{(\alpha)}) d\sigma + \\
& + \int_{\Sigma_5} s(\tilde{h}_j^{(\alpha)}\tilde{\varphi}^{*(\beta)} - \tilde{h}_j^{(\beta)}\tilde{\varphi}^{*(\alpha)})n_j d\sigma + \int_{\Sigma_6} s(\tilde{h}^{*(\alpha)}\tilde{\varphi}^{(\beta)} - \tilde{h}^{*(\beta)}\tilde{\varphi}^{(\alpha)}) d\sigma + \\
& + \frac{1}{\theta_0} \int_{\Sigma_7} (\tilde{\theta}^{*(\alpha)}\tilde{q}_j^{(\beta)} - \tilde{\theta}^{*(\beta)}\tilde{q}_j^{(\alpha)})n_j d\sigma - \frac{1}{\theta_0} \int_{\Sigma_8} (\tilde{\theta}^{(\alpha)}\tilde{q}^{*(\beta)} - \tilde{\theta}^{(\beta)}\tilde{q}^{*(\alpha)}) d\sigma = 0.
\end{aligned} \quad (22)$$

Finally, we obtain the desired equation (20) by inverting (22) and using the convolution theorem for Laplace transforms.

## References

- [1] A. C. Eringen: *Linear Theory of Micropolar Elasticity*. Int. Math. Mech. **15**, 909 (1966).
- [2] A. C. Eringen: *Foundations of Micropolar Thermoelasticity*. Springer-Verlag, Vienna (1970).
- [3] A. C. Eringen: *Theory of Micropolar Elasticity*. In Fracture (Edited by H. Leibowitz), Vol II, Academic Press, New York, 622 (1968).
- [4] A. C. Eringen and C. B. Kafadar: *Polar Field Theories*. In Continuum Physics (Edited by A. C. Eringen), Vol IV, Part I, Academic Press, New York, 1 (1976).
- [5] M. Ciarletta and D. Ieşan: *Non-Classical Elastic Solids*. In Pitman Research Notes in Mathematics Series (Edited by Longman Scientific & Technical), 293 (1993).
- [6] D. S. Chandrasekharaiah: *Heat-flux dependent micropolar thermoelasticity*, Int. J. Engng. Sci. **24**, 1389 (1986).
- [7] M. Ciarletta: *Sui processi termoelastici per continui micropolari*, Atti Sem. Mat. Fis. Univ. Modena, **39**, 103 (1991).
- [8] J. W. Nunziato and S. C. Cowin: *A non linear theory of elastic materials with voids*, Arch. Rat. Mech. Anal. **72**, 175 (1979).
- [9] S. C. Cowin and J. W. Nunziato: *Linear elastic materials with voids*, J. Elasticity **13**, 125 (1983).
- [10] M. A. Goodman and S. C. Cowin: *A continuum theory for granular materials*, Arch. Rat. Mech. Anal. **44**, 249 (1972).
- [11] F. Passarella: *Some results in micropolar thermoelasticity*, Mechanics Research Communications, **23** No. 4, 349 (1996).



- [12] M.A. Biot: *Thermoelasticity and irreversible thermodynamics*, J. Appl. Phys. **7**, 240 (1956)
- [13] H. Parkus: *Variational Principles in Thermo-and Magneto-Elasticity*, Springer-Verlag, Vienna, **11** (1972)
- [14] D. Graffi: *Sui teoremi di reciprocità nei fenomeni non stazionari*, Atti Accad. Sc. Ist. Bologna, Serie 11, **10**, 33 (1963).
- [15] D. S. Chandrasekharaiah: *Variational and reciprocal principles in micropolar thermoelasticity*, Int. J. Engng. Sci. **25**, 55 (1987).